

50

3 (Sem-3) PHY M 1

2016

PHYSICS

( Major )

Paper : 3.1

( **Mathematical Methods-III and Electrostatics** )

Full Marks : 60

Time : 3 hours

*The figures in the margin indicate full marks  
for the questions*

GROUP—A

( **Mathematical Physics** )

( Marks : 25 )

1. Answer the following questions : 1×3=3
- (a) In matrices, find the value of  $(A+B+C)^2$ .
- (b) Show that  $(A^2)^{-1} = (A^{-1})^2$ .
- (c) What is the rank of a zero matrix?

2. Check whether

$$\begin{pmatrix} i/2 & \sqrt{3}/2 \\ \sqrt{3}/2 & i/2 \end{pmatrix}$$

is a unitary matrix.

2

3. Answer any *two* of the following questions :

5×2=10

(a) (i) For an orthogonal matrix, if  $\lambda$  is an eigenvalue, what is the other value?

1

(ii) If

$$A_\alpha = \begin{pmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{pmatrix}, \quad A_\beta = \begin{pmatrix} \cos\beta & \sin\beta \\ -\sin\beta & \cos\beta \end{pmatrix}$$

check whether  $A_\alpha A_\beta = A_{\alpha+\beta}$  is correct or not.

2

(iii) If

	Room I	Room II	
$A =$	$\begin{pmatrix} 10 & 12 \\ 9 & 14 \\ 15 & 14 \end{pmatrix}$		Flat I
			Flat II
			Flat III

gives the power consumed in two rooms within three flats and

$$X = \begin{pmatrix} 10 \\ 5 \end{pmatrix} \begin{matrix} \text{Room I} \\ \text{Room II} \end{matrix}$$

gives the number of electrical items in rooms, then what information does  $Y = AX$  yield and where is its highest value?

2

(b) (i) If

$$A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}, \text{ then } A^n = \begin{pmatrix} \cos n\theta & -\sin n\theta \\ \sin n\theta & \cos n\theta \end{pmatrix}$$

What does this result mean geometrically?

1

(ii) If  $A$  and  $B$  are Hermitian matrices, show that  $AB - BA$  is skew-Hermitian whereas  $AB + BA$  is Hermitian.

2

(iii) Compute the adjoint of a matrix

$$A = \begin{pmatrix} 0 & 3 & 2 \\ -1 & 2 & 5 \\ 5 & 0 & 3 \end{pmatrix}$$

2

(c) (i) Derive the expression for the force  $F'$  acting on a body in a constant rotating frame in terms of applied force  $F$  and two other fictitious forces. Name the fictitious forces.

3+1=4

(ii) What is the effect of diurnal rotation of the earth on the acceleration due to gravity of earth at a place where latitude is  $\lambda$ ?

1

4. Answer either (a), (b) or (c), (d) : 5×2=10

(a) (i) If  $A = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$

then find the value of  $A^n$ . 2

(ii) In an electrical network

$$I_1 - I_2 + I_3 = 0$$

$$2I_2 - 3I_3 = 0$$

$$5I_1 + 3I_2 = 2$$

Find the currents by matrix method. 3

(b) (i) If

$$A = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & a & 1 \end{pmatrix}, \quad A^{-1} = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ -4 & 3 & c \\ \frac{5}{2} & -\frac{3}{2} & \frac{1}{2} \end{pmatrix}$$

then find the value of  $a+c$ . 3

(ii) If

$$A = \begin{pmatrix} 2 & -3 \\ 0 & 4 \end{pmatrix}, \quad B = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$$

find  $A-B$  and also a symmetric matrix out of it. 2

(c) (i) Verify Cayley-Hamilton theorem for the matrix

$$A = \begin{pmatrix} 1 & 1 & 0 \\ 3 & 0 & 1 \\ 2 & 3 & 1 \end{pmatrix}$$

3

(ii) If  $A = \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix}$

then using the value of

$$A^2 - 5A + 7I = 0$$

find the value of  $A^{-1}$ . 2

(d) (i) Given

$$x_1 = 3y_1 + 2y_2$$

$$x_2 = -y_1 + 4y_2$$

Find the transformation equation for  $y_1, y_2$  by matrix method. 3

(ii) If

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

satisfies the equation

$$x^2 - (a+d)x + k = 0$$

then find the relation between

$$k, a, b, c, d \quad 2$$

( 6 )

GROUP—B

( **Electrostatics** )

( Marks : 35 )

5. Choose the correct option : 1×3=3

(a) The relation  $D = \epsilon E$  is true for

(i) any medium

(ii) homogeneous medium

(iii) isotropic medium

(iv) homogeneous and isotropic medium

(b) Uniqueness of electric field strength  $E$  means


(i)  $V_1 = V_2$

(ii)  $\nabla V_1 = \nabla V_2$

(iii)  $V_1 = V_2 + \text{constant}$

(iv) Both (ii) and (iii)

- (c) A Gaussian surface in the figure below is shown by the dotted line :



The electric field on the surface will be

- (i) due to  $q_1, q_2$  only
  - (ii) due to  $q_2$  only
  - (iii) due to all
  - (iv) zero
6. Answer the following questions : 3×2=6

- (a) Given an electric field in a limited region surrounding an origin in the  $X, Y$  plane for which the potential is represented by

$$\phi = ax^2 + C$$

where  $a, C$  are positive constants. Find the components of the field intensity. Where is the potential extremum? Where is the field intensity a minimum?

$$1+1+1=3$$

Or

Show that  $K = 1 + \chi$

where  $K$  = dielectric constant

$\chi$  = susceptibility

Define polarization charges.

2+1=3

- (b) Check which of the two expressions below for electrical potential is applicable for a charged region. Correspondingly find the charge density :

2+1=3

(i)  $3x^2 + y^2 + 2z^2$

(ii)  $x^2 - y^2 + 8z$

7. Answer either (a) or (b) :

6

- (a) (i) Find the electric field at a point located at a distance  $r_1$  from the axis of a dipole of length  $d$ . Show that if  $d/r_1 \ll 1$ , the field at that behaves as  $E = 2p/r_1^3$ ,  $p$  = dipole moment.

3

- (ii) Define equipotential surface. What is the direction of electric field at a point on equipotential surface?


3



- (b) (i) A sphere of radius  $b$  is uniformly charged by charge density  $\rho$ . Calculate the electrostatic energy of the sphere. 3
- (ii) Show that the divergence of electric field of a point charge vanishes. 3

8. Answer any two questions :  $10 \times 2 = 20$

- (a) (i) An electric dipole of length 2 mm having charge of value  $q = 2.0 \times 10^{-8}$  C is placed near a long line charge of density  $4.0 \times 10^{-4}$  cm $^{-1}$  as shown in the figure



such that the negative charge is at a distance of 20 m from the line charge, the force acting on the dipole is  $0.11k$  Newton. Find  $k$ . 5

- (ii) Establish the boundary conditions satisfied by electric field  $E$  and electric displacement vector  $D$  at the boundary between two dielectrics. 5

- (b) (i) Using Laplace's equation, show that the electric field is constant in the region between the two parallel plates and it is toward the plate of lower potential.

5

- (ii) There is a solid sphere of radius  $R$  having volume charge density

$$\rho = \rho_0 (1 - r/R)$$

where  $\rho_0$  is constant and  $r$  is the distance from the centre of the sphere. Find the electric intensity  $E$  at a point inside the sphere using Gauss' law.

5

- (c) (i) Draw the field lines of  $\vec{E}$ ,  $\vec{P}$ ,  $\vec{D}$  in the region between the plates of a capacitor (of thickness  $d$ ) with the dielectric (of thickness  $t$ ) in between the plates (given  $d > t$ ). Show that  $\vec{D} = \epsilon_0 \vec{E} + \vec{P}$ .

3+2=5

- (ii) Deduce the relation between dielectric constant of a fluid and its polarizability.

5

- (d) (i) Define electrical image. Find the value of surface density of the induced charge on an infinite conducting plane due to a point charge. Draw the necessary figure. State the region where Laplace's equation is satisfied in such a case.

1+6+1=8

- (ii) An electron is distant  $10 \text{ \AA}$  from an infinite plane conductor. Calculate the force experienced by the proton. 2

★ ★ ★